


Childhood Vaccination Schedule

| | Hepatitis B | Rotavirus | Diphtheria Tetanus Pertussis Hepatitis B Poliomyelitis <i>Haemophilus influenzae</i> type b | Conjugate Pneumococcal (13vPCV) | <i>Haemophilus influenzae</i> type b Meningococcal C | Measles Mumps Rubella | Hepatitis A | Diphtheria Tetanus Pertussis | Measles Mumps Rubella Varicella | Diphtheria Tetanus Pertussis Poliomyelitis | Human Papillomavirus | Varicella | Polysaccharide Pneumococcal | Adult Diphtheria Tetanus Pertussis | Influenza REPEAT YEARLY | |
|-----------|--------------------------------------|---------------------------------------|--|---------------------------------------|---|---|---------------------------------|---|---|---|------------------------------------|--|---------------------------------------|---|---|---|
| | Engerix [®] -B 0.5ml IMI | Rotarix [®] 1.5ml ORAL | INFANRIX [®] hexa 0.5ml IMI | Prevenar 13 [®] 0.5ml IMI | Menitorix [®] 0.5ml IMI | M-M-R [®] II 0.5ml SC or [®] Priorix [®] 0.5ml IMI | VAQTA [®] 0.5ml IMI | Tripacel [®] 0.5ml IMI or [®] Infanrix [®] 0.5ml IMI | Priorix- Tetra [®] 0.5ml SC or [®] ProQuad [®] 0.5ml SC | Infanrix [®] IPV 0.5ml IMI | Gardasil [®] 0.5ml IMI | Varivax [®] 0.5ml SC or [®] Varilrix [®] 0.5ml SC | Pneumovax23 [®] 0.5ml IMI | Boostrix [®] 0.5ml IMI | Flu Quadri [™] Junior 0.25ml IMI (<3 years) | Fluarix [®] Tetra 0.5ml IMI (≥3 years) |
| Birth ● | ✓ | | | | | | | | | | | | | | | |
| 6 weeks | | ✓ | ✓ | ✓ | | | | | | | | | | | | |
| 4 months | | ✓ | ✓ | ✓ | | | | | | | | | | | | |
| 6 months | | | ✓ | ✓ | | | | | | | | | | | | ★ 6 Months |
| 12 months | | | | | ✓ | ✓ | ☐ | | | | | | | | | ★ |
| 18 months | | | | ■ | | | ☐ | ✓ | ✓ | ⚡ | | | | | | ★ |
| 4 years | | | | | | | | | | ✓ | | | | | | ★ |
| 5 years | | | | | | | | | | | | | | | | ★ |
| 12 years | | | | | | | | | | | ✓✓✓ | | | | | |
| 13 years | | | | | | | | | | | | ✓ | | ✓ | | |
| 15 years | | | | | | | | | | | | | ■ | | | ■ |



Vaccine notes:

| | | | |
|-----|---|---|---|
| ✓ | = All children. | ★ | = Indigenous children only aged 6 months to less than 5 years. All children with a chronic medical condition 6 months of age and over. Give 2 doses, 28 days apart to children under 9 years of age who are receiving influenza vaccine for the first time in their life. |
| | = ORAL VACCINE: First dose must be given by 14 weeks and 6 days of age; Second dose must be given by 24 weeks and 6 days of age. | ⚡ | = NEVER to be given as the 1 st dose of MMR containing vaccine. |
| ■ | = Indigenous people only. | ● | = BCG vaccine - for eligible groups and vaccine eligibility please refer to the CDC Immunisation Website: www.health.nt.gov.au/Centre_for_Disease_Control/Immunisation |
| ☐ | = All Indigenous children. Non-Indigenous children living in remote communities. | | |
| ✓✓✓ | = Requires 3 doses given at 0, 2 and 6 months. | | |